



SURFICIAL GEOLOGY SHEFFIELD LAKE MAP AREA (NTS 12H/07)

MAP 2016-14

SYMBOLS

Geological boundary	
Scarp face at edge of terrace	
Cirque	
Esker (flow direction known or assumed, unknown)	
Meltwater channel (small, large)	
Crestline of major moraine ridge	
Trend of ribbed or minor moraine ridges	
Beach ridges	
Sand dunes	
Avalanche track	
Drumlin (direction known, unknown)	
Crag-and-tail hill	
Till ramp	
Fluting	
Rôche moutonnée	
Striation (direction known, unknown) (numbers indicate relative age)	
Kettle hole (small, large)	
Sinkhole (small, large)	
Observation site	
Delta	
Radiocarbon age	

Note: All symbols and classifications may not occur on this map.

Geology by J.S. Organ.

GIS / digital cartography by T.J. Sears.

Digital elevation data supplied by the Shuttle Radar Topography Mission (SRTM), a partnership between NASA and the National Geospatial-Intelligence Agency (NGA). Flown aboard the NASA Space Shuttle Endeavour (11 - 22 February 2000). Additional information available from, <http://edc.usgs.gov/products/elevation/srtmbl.html>.

The age dates and glacial striations, where included, on this map have been obtained from Taylor (2001a, b).

Elevation in feet above mean sea level. Contour interval 10 metres.

Copies of this map may be obtained from the Geoscience Publications and Information Section, Geological Survey, Department of Natural Resources, Government of Newfoundland and Labrador, P.O. Box 8700, St. John's, NL, Canada, A1B 4J6. <http://edc.usgs.gov/products/elevation/srtmbl.html>.

This map is subject to review and revision. Comments to the author concerning errors or omissions are invited.

Base maps published by Surveys and Mapping Branch, Department of Natural Resources, Ottawa, Canada.

OPEN FILE 012H07/2186

This map supersedes Map 2011-08, Open File 12H07/2056.

PUBLISHED 2016.

Department <http://www.nr.gov.nl.ca/mr/>

Geological Survey: <http://www.nr.gov.nl.ca/mr/mines/geoscience/>

E-mail: pulb@nr.gov.ca

References


- Taylor, D.M. (Compiler)
2001a Carbon-14 date list for Newfoundland and Labrador. Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey. <http://gis.geosur.gov.na>
- 2001b Newfoundland and Labrador Stratigraphic Database. Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey. <http://gis.geosur.gov.na>

Recommended citation

Organ, J.S.
2016 Surficial geology of the Sheffield Lake map area (NTS 12H07). Government of Newfoundland and Labrador, Department of Natural Resources, Geological Survey, Map 2016-14, Open File 012H07/2186.

Note: Open file reports and maps issued by the Geological Survey Division of the Newfoundland and Labrador Department of Natural Resources are made available for public use without being formally edited or peer reviewed. They are based upon preliminary data and evaluation. The purchaser agrees not to provide a digital reproduction or copy of this product to a third party. Derivative products should acknowledge the source of the data.

Disclaimer: The Geological Survey, a division of the Newfoundland and Labrador Department of Natural Resources ("the publisher"), retains the sole right to copyright and all other intellectual property rights in the products it publishes. That is, these publications assume no legal liability for any derivative changes or misrepresentations found in items sold produced. That is, these publications assume no legal liability for any derivative products. Furthermore, the Geological Survey assumes no liability with respect to digital reproductions or copies of original products or for derivative products made by third parties. Please consult with the Geological Survey to ensure originally and correctness of data and/or products.



LEGEND

Each outlined area is assigned a classification consisting of up to three generic categories and modifiers that designate the types of deposits within each area. Each category within a classification is listed in order of dominance and is separated from the other categories by a slash (e.g., TvR). Generally, the areas are divided so that up to 90 percent of the area is covered by one type of deposit, and the remaining area is covered by other types. The classification also includes the percentage of landforms within the outlined area, but those that form less than 5 percent of the area are not included in the classification. Four variations of the landform system are as follows:



- Where three different landforms are included in a single map unit they are easily separated by a single slash (/) and their relative percentages are (60 - 80), (15 - 35), and (5 - 15).
- Where two landforms are included in a single map unit, a double slash (//) or single slash (/) is used to separate them, and their relative percentages are (85 - 95) // (5 - 15) for double slash, or (60 - 85) / (15 - 20) for a single slash.
- A hyphen between two landform types indicates that they are approximately equal in area. For example, Tv-Rc indicates that till veneer and rock concealed by vegetation or a thin regolith are equal in area.
- A composite symbol is used to show combinations of the above cases. For example, T indicates that about 60 - 85 percent of the area is covered by fluvial sediment, 15 - 40 percent by glaciogenic sediments, and is underlain by bedrock.

GENETIC CLASSIFICATION

Symbol	Depositional Environments	Origin and Characteristics of Materials
O	Bog	Poorly drained accumulations of peat, peat moss and other organic matter; developed in areas of poor drainage
F	Fluviatil	Alluvium consisting of silt and clay to bouldery gravel, forms terraces and plains associated with modern stream channels; their floodplains and deltas, usually less than 1 m thick, deposited by fluvial action at or below maximum flood levels
C	Cultivation	Coarse-grained bedrock-derived materials; may include sand, silt or clay, accumulates on the lower parts, or at the base of steep rock faces; transported by gravity
E	Aeolian	Medium- to fine-grained sand and silt, well sorted, poorly compacted; commonly forms as dunes up to 10 m high; transported and deposited by wind
G	Glaciogenic	Fine-grained sand to coarse-grained cobble gravel; forms plains, ridges (eskers), hummocks, terraces and deltas; generally greater than 1 m thick; deposited as outwash from ice-contact or proglacial position
L	Lacustrine	Silt, clay, gravel and sand; forms as plains and blankets; silt and clay are deposited in freshwater lakes from suspension; sand and silt by lake-floor currents; gravel and sand by shoreline wave action
M	Marine	Clay, silt, gravel and boulders; sand is present in some places, generally moderately to well sorted and commonly stratified, but may be massive; forms beach ridges, deltas, terraces and bars deposited in marine environment; gravel and sand are formed by shoreline wave action; may include shells, clay and sand deposited from storm and tsunami currents; gravel is generally a wave-washed lag
Tv	Glacial	Includes all types of till; composed of diamictite; transported and subsequently deposited by or from glacier ice with no significant sorting by water. These include relatively thin (Tv) or thicker (Tb, Te, Tp, T) till with little or no surface expression; features produced by actively flowing ice (Td, Ti, Tr) or sediment deposited through ice disintegration (Th, Tk)
Tb, Te, Tp, T		
Td, Ti, Tr		
Th, Tk		
R	Rock	Bedrock, either exposed (R) or concealed by vegetation (Rc)
Rc		

MORPHOLOGY

Symbol	Morphology	Description
a	apron	A relatively gentle slope at the foot of a steeper slope, commonly used to describe colluvium at the base of a rock escarpment; consists of materials derived from the usually steeper upper slope
b	blanket	Any deposit greater than 1.5 m thick; minor irregularities of the underlying unit are masked but the major topographic form is still evident
c	concealed by vegetation	Vegetation accumulates on either colluvial surfaces or a thin layer of angular frost-shattered and fragmented bedrock fragments overlying bedrock; includes areas of shallow (less than 1 m), discontinuous, overlying vegetation
d	drumlinoid	Elongate (ridge(s)) between 1.5 and 20 m high, and 200 m and 2000 m long; ridges elongate and end pointing in the up-ice direction and gently curving sides that taper in the down-ice direction; exhibit a convex longitudinal profile, commonly with a steeper slope in the up-ice direction; consist of subglacially deposited deposits shaped in a streamlined form parallel to the direction of glacial flow. Commonly consist of till, although some may contain stratified drift; may have a rock core; may have a rock cap
e	eroded and dissected	A series of closely spaced gullies or deeply incised channels; can have a dendritic pattern or be a single straight or arcuate channel; gullies and channels may contain underflow streams
f	fan	A gently sloping accumulation of debris carried by a stream issuing from a valley or gully; has a fan-shaped mouth at the point of entry; often contains a central channel; may result from the deposition of material as the stream swings back and forth along the lowland; fluvial fans are usually derived from eroded glacial and glaciogenic deposits; glaciogenic fans (deltas) are deposited in marine environments; cultural fans are derived from bedrock and are usually steeper (i.e., more than 10°)
h	hummock	An apparently rounded assemblage of knobs, mounds, ridges and depressions without any pronounced parallelism, significant form or orientation; formed by glacial melting during ice stagnation and deglaciation; includes subglacial, englacial, supraglacial and terrestrial hummocks
k	kettle	A basin or bowl-shaped closed depression or hollow in glacial drift; results from the melting of a buried block of bedrock or debris left by meltwater; may contain a small body of water
i	linedated	Elongate spindle-shaped ridges between 6 and 60 m high, 75 and 300 m wide and up to 4000 m long; ridges are commonly straight-sided, tapered at one or both ends, and have a flat longitudinal profile; may be subglacially formed deposits shaped in a streamlined form parallel to the direction of ice flow; common in complex terrain; although some may contain stratified drift, may have a rock core; includes slope linedated bogs (O)
p	plain	A comparatively flat, level, or slightly undulating tract of land; materials are either till, glaciogenic, alluvium, marine, lacustrine or organic sediments; bedrock features are commonly masked by the overlying material
r	ridge	Narrow, elongated and commonly steep-sided feature that rises above the surrounding terrain; materials are either rock, till, glaciogenic, fluvial, marine, lacustrine, aeolian, or organic sediments; includes string bogs (O)
t	terrace	Long, narrow, level or gently inclined step-like surface, bounded along one edge by a steeper descending slope or scarp and along the other by a steeper ascending slope or scarp; materials are either till, glaciogenic, or lacustrine sediments; generally formed by fluvial and glaciogenic erosion or marine wave action
v	veener	Any deposit less than 1.5 m thick; morphology of the underlying unit is evident
w	weathered	A thin layer, generally less than 1 m thick, of frost-heaved and frost-shattered bedrock fragments
x	weathered complex	Commonly used to indicate numerous weathered ridges that are closely spaced; can be used when any genetic category exhibits numerous surface expressions in a small area, and in which no single element can be defined at this scale



Digital Elevation Model (DEM) from Shuttle Radar Topography Mission (SRTM) data of the Sheffield Lake map area (shaded from the northeast). The image provides surface information not readily illustrated on the surficial map. Terrain variability is evident, with areas of bedrock highlighted as rougher textured areas (consistent with surficial map), and areas of thicker till shown as smoother textured areas.